READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CLIPSHIELD®

LIQUID BLOWFLY TREATMENT FOR SHEEP

ACTIVE CONSTITUENT: 500 g/L CYROMAZINE

Protects long wool sheep for up to 14 weeks from strike by blowfly (*Lucilia cuprina*), including organophosphate resistant strains when applied by jetting or dipping.

(covine)®

DIRECTIONS FOR USE:

Restraints:
DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Contraindications:
This product is contraindicated for use on open wounds e.g. those resulting from marking or mulesing operations.
This product is contraindicated for use on sheep with less than 6 weeks wool as a reduced period of protection may result.
Dipping sheep heavily infested with grass seeds is not recommended.
Heavy rain following the application of this product could diminish the period of protection.
If used in jetting races a reduced period of protection may result.

Dosage and Administration:

PEST	TREATMENT	INITIAL CHARGE	TOPPING UP	CRITICAL COMMENTS
Sheep blowfly control (*Lucilia cuprina*)	Jetting	2 L per 1000 L	–	1. Jet at onset of blowfly wave. 2. Apply 2 to 4L of jetting fluid per sheep, depending on sheep size and wool length. 3. Ensure saturation to the skin 4. Pay particular attention to treatment of the crutch, pizzles and polls. 5. This product is not recommended for use in jetting races because a reduced period of protection may result.
	Dipping (plunge and shower and constant replenishment)		2 L per 1000 L	1. Dip sheep at onset of blowfly wave. 2. Thoroughly saturate sheep. When using a shower dip, run top and bottom sprays independently so as to ensure maximum pressure and therefore penetration. 3. The product is a non-stripping dip so there is no need to reinforce during dipping.
	Dressing		–	1. Clip the wool surrounding the strike, treat the struck area and the surrounding wool with the Relyon (Australia) Pty Ltd recommended diazinon knockdown product, then thoroughly jet the sheep with Clipshield Liquid Blowfly Treatment For Sheep. 2. This product is a slow killing larvicide and when used on its own, larvae may take 3 to 4 days to die.

General Directions:

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

This product is designed for use on sheep with 6 weeks wool growth (long wool). Dipping sheep heavily infested with grass seeds is not recommended. Heavy rain following the application of this product could diminish the period of protection. If used in jetting races a reduced period of protection may result.

MIXING. Spray preparation: SHAKE WELL BEFORE USE
Clipshield Liquid Blowfly Treatment For Sheep may settle after storage for some weeks. Stir product or invert the container several times before opening. Pour the liquid into the vat through a strainer to remove any particles or dry flakes, which may occur in hot storage conditions.
Product may be blended with water and applied as spray. Partly (30%) fill the tank with water. Put the required amount of Clipshield Liquid Blowfly Treatment For Sheep for the area to be sprayed into water containers. Add this correct amount of pre-mixed Clipshield Liquid Blowfly Treatment For Sheep into the tank with the agitation system operating. Fill up to the correct volume with water. The product is readily soluble in both soft and hard water. Tank-mixtures should be tested for physical compatibility and efficacy before use. If tank-mixing, Clipshield Liquid Blowfly Treatment For Sheep must be in suspension before adding other insecticides or surfactant. Wash equipment thoroughly after use.

DILUTION RATE:
Dilution rate 2 mL per 1 litre of water
E.g. 400 mL per 200 litres or 2 L per 1000 L

MAINTENANCE OF DIPS
Clean out dip, sump and spray lines thoroughly before charging. Disinfect with a registered bacteriostat at the manufacturer's recommended rate.DO NOT use old fouled wash. Shower dips should be emptied and cleaned out at the end of each day's dipping. Plunge dips should be emptied and cleaned out within 5 days of charging and more frequently if conditions favour post-dipping lameness infection, except in WA where plunge dips and sumps should be emptied daily. Before use agitate well.
Jetting: Jet the sheep at onset of fly wave with efficient equipment that is able to maintain the pressure of about 550 kPa. Use 300 to 400 kPa for lambs and ewes in lamb.
Bacteriostat: This product does not contain a bacteriostat. For control of post-dipping infections a registered bacteriostat should be added at the manufacturer's recommended rate.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 7 days before slaughter for human consumption.

MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

WITHHOLDING PERIOD - WOOL: DO NOT USE less than 2 months before shearing or fibre collection.

TRADE ADVICE – EXPORT SLAUGHTER INTERVAL (ESI): DO NOT use less than 21 days before slaughter for export. Before using this product, confirm the current ESI from Relyon (Australia) Pty Ltd on (02) 9952 6666 or the APVMA website (www.apvma.gov.au/residues).

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container and preparing the spray/dip/dressing wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC gloves. When using the prepared spray/dip/dressing wear protective waterproof clothing and PVC or rubber apron, elbow-length PVC gloves, face shield and water resistant footwear. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

ADDITIONAL USER SAFETY INFORMATION:
Additional information is listed in the safety data sheet.

Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.

DISPOSAL: Triple-rinse container into the mediated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: Keep out of reach of children. Store below 30ºC (room temperature). Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Do not re-use the container or dispose of undiluted chemicals on-site. Shake empty into spray tank.

IN A TRANSPORT EMERGENCY
DIAL: 000
POLICE OR FIRE BRIGADE

APVMA Approval No: 86790/116577

(covine)®

drumMUSTER eligible container

Relyon (Australia) Pty Ltd
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW 2113
ABN 34 607 057 232 ACN 607 057 232 Phone (02) 9952 6555
covine.com.au

COVINE® is a registered trademark of Relyon (Australia) Pty Ltd.
CLIPSHIELD® is a registered Trademark of Relyon (Australia) Pty Ltd.