

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

COVINE

DELTASHIELD

POUR-ON LICE AND FLY TREATMENT FOR CATTLE

ACTIVE CONSTITUENT: 15g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from the Biting Midge (*Culicoides brevitarsus*) for at least 14 days with effective control of blood feeding *C. brevitarsus* for 6 days post treatment.



(covine)[®]

5L



COVINE DELTASHIELD POUR-ON LICE AND FLY TREATMENT FOR CATTLE**DIRECTIONS FOR USE:****Restraints: DO NOT USE on horses that may be used for human consumption.****Contraindications:** This product is contraindicated for use on animals severely stressed or in ill health. This product should not be used on horse's head forward of the ears. This product should not be applied as a single spot.**Precautions:** Facial skin contact may cause temporary tingling/burning.**Dosage and Administration:****DOSE RATE FOR CATTLE:** 5 mL per 100 kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Lignognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5 mL 10 mL 15 mL 20 mL	Up to 100 kg 101 – 200 kg 201 – 300 kg 301 – 400 kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>) Biting Midge (<i>Culicoides</i> sp.)	25 mL 30 mL For cattle over 600 kg apply 5 mL per 100 kg liveweight	401 – 500 kg 501 – 600 kg 	Apply to the whole herd – retreatment may be necessary.

DOSE RATE FOR HORSES: 5 mL per 100 kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Buffalo Fly (<i>Haematobia irritans exigua</i>)	5 mL 10 mL 15 mL	Up to 100 kg 101 – 200kg 201 – 300kg	Using the correct measured dose, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump.
Stable Fly (<i>Stomoxys calcitrans</i>)	20 mL 25 mL 30 mL For horses over 600 kg apply 5 mL per 100 kg liveweight	301 – 400kg 401 – 500kg 501 – 600kg 	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**General Directions:****Cattle:** To avoid operator insecticide contact when concurrent treatments are carried out, DELTASHIELD POUR-ON should be applied last. Operators must wear gloves.

In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product.

Apply with recommended applicator.

The efficacy of this product was not affected by moderate rainfall when tested against cattle lice.

Horses: Treatment with DELTASHIELD POUR-ON will alleviate fly annoyance related behaviour including skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated with deltamethrin at 5 times the recommended dose rate. It is recommended that the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label.**WITHHOLDING PERIODS:****Cattle: MEAT: Zero (0) days; MILK: Zero (0) days****MEAT WITHHOLDING PERIOD (HORSES): DO NOT USE on horses that may be used for human consumption.****TRADE ADVICE – EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product, confirm the current ESI from Relyon (Australia) Pty Ltd on (02) 9952 6666 or the APVMA website (www.apvma.gov.au/residues).****SAFETY DIRECTIONS:** Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.**ADDITIONAL USER SAFETY INFORMATION:** AVOID CONTACT WITH FOOD. Additional information is listed in the safety data sheet.**ENVIRONMENTAL PROTECTION:** DANGEROUS TO FISH. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.**DISPOSAL:** Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.**STORAGE:** Store below 30°C (room temperature) in original container, tightly closed in a safe place.**CONDITIONS OF SALE:** Any provision or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Ruralco Holdings Ltd and/or its associated entities shall not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.**Harmful if swallowed.**

Wear protective gloves, protective clothing and eye or face protection.

Take off protective clothing and wash before reuse.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Call a POISON CENTRE or doctor if you feel unwell.

If skin irritation occurs: Get medical advice.

Collect spillage.

Store in a well ventilated, dry place. Protect from sunlight.

RELYON®Relyon (Australia) Pty Ltd
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW 2113
ABN 34 607 057 232 ACN 607 057 232 Phone (02) 9952 6666
covine.com.au©RELYON is a Registered Trademark of Ruralco Holdings Limited
COVINE® is a registered trademark of Relyon (Australia) Pty Ltd.
DELTASHIELD™ is a trademark of Relyon (Australia) Pty Ltd.