

COVINE

# TOROME<sup>TM</sup> LV

POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 10 g/L IVERMECTIN

For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle.



(covine)<sup>®</sup>

**Indications:**

For the treatment and control of adult and some immature stages of:

**Gastro-intestinal roundworms**

Barber's Pole Worm (*Haemonchus placei*)

Hookworm (*Bunostomum phlebotomum*)

Intestinal Threadworm (*Strongyloides papillosus*)

Large Bowel Worm (*Oesophagostomum radiatum*)

Small Intestinal Worm (*Cooperia oncophora*) (*Cooperia punctata*)

Small Brown Stomach Worm (*Ostertagia ostertagi*)

(Including inhibited stages)

Stomach Hair Worm (*Trichostrongylus axei*)

Thin Necked Intestinal Worm (*Nematodirus spathiger*)

Whipworm (*Trichuris* spp.)

**Lungworms** (*Dictyocaulus viviparus*)**Sucking and Biting Lice** Biting lice (*Bovicola bovis*)

Little sucking lice (*Solenopotes capillatus*)

Long nosed sucking lice (*Linognathus vituli*)

Short nosed sucking lice (*Haematopinus eurysternus*)

**Mites** (*Chorioptes bovis*) (*Sarcoptes scabiei var bovis*)

**Buffalo Fly** (*Haematobia irritans exigua*)

**Cattle Tick** (*Boophilus microplus*)

Maximum tick efficacy is achieved 4-5 days after treatment. In planning a strategic tick control program, it should be noted that viable eggs may have been laid by engorged female ticks that drop from cattle in the initial days after treatment.

COVINE TOROMEC LV POUR-ON FOR CATTLE should be alternated with conventional dips or Pour-Ons when the program requires repeat treatments.

**Sustained Activity:** COVINE TOROMEC LV POUR-ON FOR CATTLE applied at the recommended dose of 1 mL per 20 kg bodyweight effectively controls infection with *Ostertagia* spp. (Small brown stomach worm) acquired up to 14 days after treatment, with *Dictyocaulus viviparus* (Lungworm) acquired up to 28 days after treatment and *Haematobia irritans exigua* (Buffalo Fly) up to 28 days after treatment.

**DIRECTIONS FOR USE****Dosage and Administration:**

COVINE TOROMEC LV POUR-ON FOR CATTLE is for external application to cattle only.

The recommended dose rate for COVINE TOROMEC LV POUR-ON FOR CATTLE is 1 mL/20 kg body weight. Apply along the topline of the animal.

Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg.

A representative sample of animals should be weighed before treatment either with scales or a weighband.

Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not under dose.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Liveweight (kg)	Dose Volume (mL)	No. head treated per pack
		2.5 L
51 – 100	5	500
101 – 150	7.5	333
151 – 200	10	250
201 – 250	12.5	200
251 – 300	15	166
301 – 350	17.5	142
351 – 400	20	125
401 – 450	22.5	111
451 – 500	25	100
501 – 550	27.5	90
551 – 600	30	83
601 – 650	32.5	76

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**General directions:** Rainfall on the herd within 2 hours of application may result in reduced efficacy of the product. Cattle should not be treated with COVINE TOROMEC LV POUR-ON FOR CATTLE when the hide or hair is wet since the efficacy of the product may be reduced. Temporary hair loss occasionally occurs in the application area but will resolve without further treatment.

Do not retreat with COVINE TOROMEC LV POUR-ON FOR CATTLE within the meat withholding period of 42 days.

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE less than 42 days before slaughter for human consumption.**

**MILK: Zero (0) days.**

**TRADE ADVICE – EXPORT SLAUGHTER INTERVAL**

**(ESI): DO NOT use less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Relyon (Australia) Pty Ltd on (02) 9952 6666 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**